

VOERMOL PRODUCTION LICK

Class: Protein, Energy, Mineral and Trace Mineral Supplement for Ruminants

VOERMOL PRODUCTION LICK is a complete, ready-mixed lick that provides protein, energy, minerals, vitamins and trace minerals for the optimum production of the grazing animal:

- contains molasses as energy sources;
- helps to prevent excessive mass loss in cows after calving;
- helps to prevent pregnancy toxæmia in ewes during late gestation and can also be used for flush feeding of ewes;
- ensures weight gain of young growing animals on good quality veld;
- part of the trace minerals are supplied in organic form which is highly absorbable and promotes optimal reproduction and growth;
- contains high levels of vitamin A, D3 and E. Vitamin E in combination with selenium is a powerful anti-oxidant which strengthens the animal's immunity. Vitamin D3 helps with the absorption of phosphorus and calcium;
- contains medicaments which promote energy usage from pastures.



COMPOSITION OF VOERMOL PRODUCTION LICK

REG NO V 10108 (Act 36/1947)

(g/kg)

(mg/kg)

Crude Protein* (min)	250	Manganese	300
Urea (max)	55	Copper	50
Crude Fibre (max)	100	Cobalt	0.98
Moisture (max)	160	Iron	150
Calcium (min/max)	10/12	Iodine	4
Phosphorus (min)	8	Zinc	290
Potassium (min)	18	Selenium	0.9
Magnesium (min)	5	Vitamin A (IU/kg)	10 000
Sulphur (min)	4	Vitamin D3 (IU/kg)	2 000
		Vitamin E (mg/kg)	50
		Energy est. (MJ ME/kg)	8
*67.4 % derived from NPN		Contains medicaments	

Packaging – 50 kg

See urea warning

NB. Contains medicaments and must therefore not be fed to horses.
MEDICATION WARNING: This product contains registered feed additives. Feed strictly according to the prescribed instructions.

FEEDING INSTRUCTIONS

Use **Voermol Production Lick** as a protein and energy supplement to hay and/or veld. In consultation with a veterinarian, follow a complete dosing and vaccination programme inclusive of *Clostridium spp.*

RECOMMENDED INTAKE

(g/animal/day)

Cattle: 1 000 – 1 500*

Sheep: 200 – 270*

*Maximum allowable intake of lick.



UREA WARNING

UREA

The majority of licks contain urea that can be dangerous under certain conditions. During the formulation and manufacturing process of **Voermol licks**, great care is taken to minimise the risks of feeding urea. The possibilities of urea poisoning can be further reduced by taking note of the following:

- use licks as prescribed;
- limit lick intake to the prescribed maximum levels;
- feed the licks together with sufficient grazing and/or roughage;
- take special care during feed scarcity (drought conditions) to prevent high lick intake;
- prevent excessive lick intake in the initial stages (salt hunger) by allowing the animals access to **Voermol Rumevite 6P** for the first 14 days;
- do not let licks get wet.

See urea warning on the product packaging.

UREA WARNING

This farm feed contains NPN (Non-Protein Nitrogen) sources and must therefore be fed strictly according to instructions for use.

- Vinegar is an effective remedy against NPN poisoning. Mix with an equal amount of water. Dose half a bottle per calf/sheep or 2 – 4 bottles per head of cattle (1 bottle = 750 ml).
- Protect this farm feed against rain. NPN is soluble and animals drinking such a solution could be poisoned.
- Do not feed this farm feed indiscriminately with other NPN containing farm feeds. Consult an animal scientist.
- Mix concentrate thoroughly with the prescribed ingredients.
- This is a supplement and not a feed. Sufficient grazing and/or roughage must be available at all times.
- Keep lick troughs filled and prevent gluttonous eating by hungry animals. A constant daily intake can help prevent poisoning.
- Before feeding a NPN containing lick, feed an ordinary salt / phosphate lick for at least 7 days.

GMO STATEMENT

These animal feeds may contain ingredients derived from GMO (genetically modified organisms) origin.

