


PRODUCTION SUPPLEMENTS

VOERMOL SUPERMOL

Class: Protein, Mineral and Energy Supplement for Ruminants

VOERMOL SUPERMOL is a protein, energy, mineral supplement for ruminants on green veld pastures:

- supplies energy and protein, which are often the limiting nutrients on green grazing as well as in certain non-fertilised, cultivated pastures;
- provides phosphorus and trace minerals, the first limiting nutrients in green grazing;
- ensures adequate intake because it is molasses based and thus palatable;
- is not dusty and thus not susceptible to being blown away by the wind;
- contains medicaments which promote energy usage from green veld. Medicaments also reduce the risk of ketosis, pregnancy toxæmia and coccidiosis, as well as abortions in Angora ewes.



COMPOSITION OF VOERMOL SUPERMOL

REG NO V 7267 (Act 36/1947)

(g/kg)

(mg/kg)

Crude Protein* (min)	200	Manganese	240
Urea (max)	38	Copper	80
Crude Fibre (max)	100	Cobalt	1
Moisture (max)	160	Iron	75
Calcium (min/max)	8/12	Iodine	5
Phosphorus (min)	6	Zinc	350
Magnesium (min)	6	Selenium	1
Sulphur (min)	3.5	Energy estimate (MJ ME/kg)	9
*60.5 % derived from NPN		Contains medicaments	

Packaging – 50 kg

See urea warning

FEEDING INSTRUCTIONS

Feed in lick troughs. Intake can be controlled to the desired level by adding 10 % salt. Feed to grazing animals on green veld or non-fertilised cultivated pastures.

RECOMMENDED INTAKE

(g/animal/day)

Cattle:

1 000 – 1 200

Sheep:

180 – 200



VOERMOL

If the quality of grass decreases, mix the following mixture and feed 2.4 kg/animal/day and 400 g/sheep/day.

Mixture		(kg)
Voermol Supermol		50
Milled grain / Hominy Chop		50
Total		100
COMPOSITION OF MIXTURE (g/kg)		
Crude Protein*	(min)	115
Urea	(max)	15
Moisture	(max)	140
Calcium	(max)	4
Phosphorus	(min)	4
Energy estimate	(MJ ME/kg)	10.5
*37.4 % derived from NPN		

NB. Voermol Supermol contains medicaments and must therefore not be fed to horses.



UREA WARNING

UREA

The majority of licks contain urea that can be dangerous under certain conditions. During the formulation and manufacturing process of **Voermol** licks, great care is taken to minimise the risks of feeding urea. The possibilities of urea poisoning can be further reduced by taking note of the following:

- use licks as prescribed;
- limit lick intake to the prescribed maximum levels;
- feed the licks together with sufficient grazing and/or roughage;
- take special care during feed scarcity (drought conditions) to prevent high lick intake;
- prevent excessive lick intake in the initial stages (salt hunger) by allowing the animals access to **Voermol Rumevite 6P** for the first 14 days;
- do not let licks get wet.

See urea warning on the back of the bag.

UREA WARNING

This farm feed contains NPN (Non-Protein Nitrogen) sources and must therefore be fed strictly according to instructions for use.

- Vinegar is an effective remedy against NPN poisoning. Mix with an equal amount of water. Dose half a bottle per calf/sheep or 2 – 4 bottles per head of cattle (1 bottle = 750 ml).
- Protect this farm feed against rain. NPN is soluble and animals drinking such a solution could be poisoned.
- Do not feed this farm feed indiscriminately with other NPN containing farm feeds. Consult an animal scientist.
- Mix concentrate thoroughly with the prescribed ingredients.
- This is a supplement and not a feed. Sufficient grazing and/or roughage must be available at all times.
- Keep lick troughs filled and prevent gluttonous eating by hungry animals. A constant daily intake can help prevent poisoning.
- Before feeding a NPN containing lick, feed an ordinary salt / phosphate lick for at least 7 days.

GMO STATEMENT

These animal feeds may contain ingredients derived from GMO (genetically modified organisms) origin.



VOERMOL