

PROTEIN LICKS

VOERMOL EKONOLICK

Class: Protein, Mineral and Trace Mineral Supplement for Ruminants

VOERMOL EKONOLICK is a ready-mixed, energy rich, protein lick with a high phosphorus content for cattle, sheep and goats on dry pastures:

- is molasses based, not dusty and will not blow away;
- as a protein source it will stimulate the appetite in order to make better use of less palatable dry veld;
- will help prevent weight loss during the dry season and therefore contribute to the increase in lamb and calf percentages;
- can be fed to cattle, sheep and goats in the same camp.



COMPOSITION OF VOERMOL EKONOLICK

REG NO V11147 (Act 36/1947) N-FF 0628

(g/kg)

(mg/kg)

Crude Protein*	(min)	240	Manganese	120
Urea	(max)	68	Copper	30
Moisture	(max)	160	Cobalt	0.3
Crude Fibre	(max)	100	Iron	150
Calcium	(min/max)	20/36	Iodine	3
Phosphorus	(min)	10	Zinc	120
Magnesium	(min)	3	Selenium	0.3
Sulphur	(min)	9	Vitamin A (IU/kg)	12 750
*88.4 % derived from NPN				

Packaging – 50 kg

See urea warning

FEEDING INSTRUCTIONS

Voermol Ekonolick is a ready-mixed lick. Feed in lick troughs to grazing animals. The intake of the lick will be influenced by the quality and quantity of the dry veld and the condition of the animals, as well as the salt content of the drinking water and the pasture.

Voermol Ekonolick is suitable for ruminant game.

RECOMMENDED INTAKE

(g/animal/day)

Cattle: 630 – 800

Sheep and Goats: 105 – 180



VOERMOL

UREA WARNING

UREA

The majority of licks contain urea that can be dangerous under certain conditions. During the formulation and manufacturing process of **Voermol** licks, great care is taken to minimise the risks of feeding urea. The possibilities of urea poisoning can be further reduced by taking note of the following:

- use licks as prescribed;
- limit lick intake to the prescribed maximum levels;
- feed the licks together with sufficient grazing and/or roughage;
- take special care during feed scarcity (drought conditions) to prevent high lick intake;
- prevent excessive lick intake in the initial stages (salt hunger) by allowing the animals access to **Voermol Rumevite 6P** for the first 14 days;
- do not let licks get wet.

See urea warning on the back of the bag.

UREA WARNING

This farm feed contains NPN (Non-Protein Nitrogen) sources and must therefore be fed strictly according to instructions for use.

- Vinegar is an effective remedy against NPN poisoning. Mix with an equal amount of water. Dose half a bottle per calf/sheep or 2 – 4 bottles per head of cattle (1 bottle = 750 ml).
- Protect this farm feed against rain. NPN is soluble and animals drinking such a solution could be poisoned.
- Do not feed this farm feed indiscriminately with other NPN containing farm feeds. Consult an animal scientist.
- Mix concentrate thoroughly with the prescribed ingredients.
- This is a supplement and not a feed. Sufficient grazing and/or roughage must be available at all times.
- Keep lick troughs filled and prevent gluttonous eating by hungry animals. A constant daily intake can help prevent poisoning.
- Before feeding a NPN containing lick, feed an ordinary salt / phosphate lick for at least 7 days.

GMO STATEMENT

These animal feeds may contain ingredients derived from GMO (genetically modified organisms) origin.



VOERMOL