


## PROTEIN LICKS

### RUMEVITE CATTLE BLOCK

Class: Protein and Mineral Supplement for Cattle

**RUMEVITE CATTLE BLOCK** is the easiest way of feeding maintenance supplements to all types of cattle;

- is ready to use and no further additives are needed;
- is a concentrated source of protein in block form which leads to a low cost per unit protein;
- it helps with digestion of dry roughage and pasture since it provides protein and trace minerals that stimulate the rumen bacteria.



#### COMPOSITION OF RUMEVITE CATTLE BLOCK

REG NO V10932 (Act 36/1947) N-FF 0627

(g/kg)		(mg/kg)	
Crude Protein* (min)	400	Manganese	160
Urea (max)	124	Copper	40
Crude Fibre (max)	20	Cobalt	0.4
Moisture (max)	140	Iron	200
Calcium (max)	48	Iodine	8
Phosphorus (min)	4	Zinc	160
Magnesium (min)	8	Selenium	0.4
Sulphur (min)	5	Vitamin A (IU/kg)	20 400

\*89 % derived from NPN

Packaging – 25 kg

**See urea warning**

#### FEEDING INSTRUCTIONS

Cattle: Supply 1 **Rumevite Cattle Block** per 15 animals.

#### RECOMMENDED DAILY INTAKE

Cattle: 375 - 600 g/animal/day.



**VOERMOL**

## UREA WARNING

### UREA

The majority of licks contain urea that can be dangerous under certain conditions. During the formulation and manufacturing process of **Voermol** licks, great care is taken to minimise the risks of feeding urea. The possibilities of urea poisoning can be further reduced by taking note of the following:

- use licks as prescribed;
- limit lick intake to the prescribed maximum levels;
- feed the licks together with sufficient grazing and/or roughage;
- take special care during feed scarcity (drought conditions) to prevent high lick intake;
- prevent excessive lick intake in the initial stages (salt hunger) by allowing the animals access to **Voermol Rumevite 6P** for the first 14 days;
- do not let licks get wet.

**See urea warning on the back of the bag.**

### UREA WARNING

This farm feed contains NPN (Non-Protein Nitrogen) sources and must therefore be fed strictly according to instructions for use.

- Vinegar is an effective remedy against NPN poisoning. Mix with an equal amount of water. Dose half a bottle per calf/sheep or 2 – 4 bottles per head of cattle (1 bottle = 750 ml).
- Protect this farm feed against rain. NPN is soluble and animals drinking such a solution could be poisoned.
- Do not feed this farm feed indiscriminately with other NPN containing farm feeds. Consult an animal scientist.
- Mix concentrate thoroughly with the prescribed ingredients.
- This is a supplement and not a feed. Sufficient grazing and/or roughage must be available at all times.
- Keep lick troughs filled and prevent gluttonous eating by hungry animals. A constant daily intake can help prevent poisoning.
- Before feeding a NPN containing lick, feed an ordinary salt / phosphate lick for at least 7 days.

### GMO STATEMENT

These animal feeds may contain ingredients derived from GMO (genetically modified organisms) origin.



**VOERMOL**