

FATTENING PRODUCTS

VOERMOL SHEEP FATTENING PELLETS

Class: Sheep Finisher Feed

VOERMOL SHEEP FATTENING PELLETS is a complete feed for the feedlot fattening of lambs:

- contains the necessary minerals, trace minerals as well as Vit. A, to achieve optimum performance;
- contains an ionophore and other medicaments to improve performance and efficiency.



COMPOSITION OF VOERMOL SHEEP FATTENING PELLETS

REG NO V12796 (Act 36/1947)

(g/kg)

(mg/kg)

Protein*	(min)	140	Manganese	40
Urea	(max)	10	Copper	9
Moisture	(max)	120	Cobalt	0.15
Crude Fibre	(max)	150	Iron	75
Fat	(min)	25	Iodine	0.6
Calcium	(max)	10	Zinc	50
Phosphorus	(min)	3	Selenium	0.16
Magnesium	(min)	2	Vitamin A (IU/kg)	4000
Sulphur	(min)	1.5	Energy (MJ ME/kg)	10.2
*28.1 % derived from NPN			Contains medicaments	

Packaging 50 kg

See urea warning

FEEDING INSTRUCTIONS

Voermol Sheep Fattening Pellets is a complete feed and is fed *ad lib*. Feed additional hay during the first 14 days of adaptation.



VOERMOL

UREA WARNING

UREA

The majority of licks contain urea that can be dangerous under certain conditions. During the formulation and manufacturing process of **Voermol** licks, great care is taken to minimise the risks of feeding urea. The possibilities of urea poisoning can be further reduced by taking note of the following:

- use licks as prescribed;
- limit lick intake to the prescribed maximum levels;
- feed the licks together with sufficient grazing and/or roughage;
- take special care during feed scarcity (drought conditions) to prevent high lick intake;
- prevent excessive lick intake in the initial stages (salt hunger) by allowing the animals access to **Voermol Rumevite 6P** for the first 14 days;
- do not let licks get wet.

See urea warning on the back of the bag.

UREA WARNING

This farm feed contains NPN (Non-Protein Nitrogen) sources and must therefore be fed strictly according to instructions for use.

- Vinegar is an effective remedy against NPN poisoning. Mix with an equal amount of water. Dose half a bottle per calf/sheep or 2 – 4 bottles per head of cattle (1 bottle = 750 ml).
- Protect this farm feed against rain. NPN is soluble and animals drinking such a solution could be poisoned.
- Do not feed this farm feed indiscriminately with other NPN containing farm feeds. Consult an animal scientist.
- Mix concentrate thoroughly with the prescribed ingredients.
- This is a supplement and not a feed. Sufficient grazing and/or roughage must be available at all times.
- Keep lick troughs filled and prevent gluttonous eating by hungry animals. A constant daily intake can help prevent poisoning.
- Before feeding a NPN containing lick, feed an ordinary salt / phosphate lick for at least 7 days.

GMO STATEMENT

These animal feeds may contain ingredients derived from GMO (genetically modified organisms) origin.



VOERMOL