

# FATTENING PRODUCTS

## VOERMOL ZILMOL

Class: Molasses and Roughage Supplement for Cattle

**VOERMOL ZILMOL** is an additive containing product for the use in feedlot finishing rations:

- contains an  $\beta$ -agonist, Zilphaterol HCl. Zilphaterol HCl increases the deposition of muscle while it suppresses the deposition of fat;
- may increase the average daily gain by about 8 %, the feed conversion ratio by about 12 % and the dressing percentage by about 3 %;
- will therefore increase the carcass weight.

By using **Voermol Zilmol** the feeding period will be increased by about 20 days. This may increase feed margin and profitability.

***Voermol Zilmol should only be fed to slaughter cattle.  
It is only registered for use in cattle feeding.***



COMPOSITION OF VOERMOL ZILMOL		
REG NO V16978 (Act 36/1947)		
		(g/kg)
Crude Protein	(min)	38
Crude Fibre	(max)	120
Moisture	(max)	160
Fat	(min)	4.5
Calcium	(max)	6
Phosphorus	(min)	0.6
Energy	(MJ ME/kg)	9.6
		Contains medicaments

Packaging – 40 kg

***NB. Voermol Zilmol contains feed additives.  
Feed strictly according to the feeding instructions.***

### FEEDING INSTRUCTIONS

- Feed Voermol Zilmol rations for the last 30 to 35 days of the fattening period;
- Refer to mixing instructions with **Voermol Beefmaker**, **Voermol SB 100** or **Voermol Super 80** for rations;
- Start feeding when oxen reach an A2 grading. Heifers can be fed the finisher feed at an earlier age;
- Allow a withdrawal period of between three and seven days before slaughter.



**VOERMOL**