

## MAINTENANCE PELLETS

### VOERMOL RAM, LAMB AND EWE PELLETS

Class: Complete Ewe and Lamb Feed

**VOERMOL RAM, LAMB AND EWE PELLETS** is a completely balanced feed pellet for the feeding of all sheep and goats:

- can be fed as is and is completely balanced;
- is in pellet form which makes it easy to feed with no milling and mixing costs;
- is precisely mixed, promotes optimum feed utilisation and eliminates wastage;
- is molasses based and therefore palatable with good intakes;
- can be fed as a supplementary feed to rams, lambs and ewes on the veld;
- is an ideal supplementary feed for sheep and goats that are kept in an enclosure or pen overnight;
- contains anionic salts and can therefore safely be fed to rams;
- contains sufficient balanced amounts of protein, energy, minerals and trace minerals for general use in sheep and goat nutrition;
- promotes good production.



#### COMPOSITION OF VOERMOL RAM, LAMB AND EWE PELLETS

REG NO V26842 (Act 36/1947)

(g/kg)

(mg/kg)

Crude Protein* (min)	130	Manganese	18
Urea (max)	14.7	Copper	6.3
Crude Fibre (max)	200	Cobalt	0.1
Moisture (max)	120	Iron	27
Calcium (min/max)	6/15	Iodine	0.1
Phosphorus (min/max)	3/3.6	Zinc	18
Magnesium (min)	0.9	Selenium	0.1
Sulphur (min)	1.35	Vitamin A (IU/kg)	3 000
*40 % derived from NPN		Energy (MJ ME/kg)	9.2

Packaging – 50 kg

See urea warning

#### FEEDING INSTRUCTIONS

**Voermol Ram, Lamb and Ewe Pellets** is a complete balanced pellet and can be given to sheep and goats as is on veld or in a small camp or pen.

#### RECOMMENDED INTAKE

As a supplementary feed to veld and hay, feed the following amounts:

Rams: 400 – 600 g/ram/day

Ewes: 300 – 500 g/ewe/day

Lambs: 200 – 300 g/lamb/day

If grazing conditions are very poor, it can be fed as a maintenance feed at 2 % to 2.5 % of bodyweight to sheep and goats.



**VOERMOL**



## UREA WARNING

### UREA

The majority of licks contain urea that can be dangerous under certain conditions. During the formulation and manufacturing process of **Voermol** licks, great care is taken to minimise the risks of feeding urea. The possibilities of urea poisoning can be further reduced by taking note of the following:

- use licks as prescribed;
- limit lick intake to the prescribed maximum levels;
- feed the licks together with sufficient grazing and/or roughage;
- take special care during feed scarcity (drought conditions) to prevent high lick intake;
- prevent excessive lick intake in the initial stages (salt hunger) by allowing the animals access to **Voermol Rumevite 6P** for the first 14 days;
- do not let licks get wet.

**See urea warning on the back of the bag.**

### UREA WARNING

This farm feed contains NPN (Non-Protein Nitrogen) sources and must therefore be fed strictly according to instructions for use.

- Vinegar is an effective remedy against NPN poisoning. Mix with an equal amount of water. Dose half a bottle per calf/sheep or 2 – 4 bottles per head of cattle (1 bottle = 750 ml).
- Protect this farm feed against rain. NPN is soluble and animals drinking such a solution could be poisoned.
- Do not feed this farm feed indiscriminately with other NPN containing farm feeds. Consult an animal scientist.
- Mix concentrate thoroughly with the prescribed ingredients.
- This is a supplement and not a feed. Sufficient grazing and/or roughage must be available at all times.
- Keep lick troughs filled and prevent gluttonous eating by hungry animals. A constant daily intake can help prevent poisoning.
- Before feeding a NPN containing lick, feed an ordinary salt / phosphate lick for at least 7 days.

### GMO STATEMENT

These animal feeds may contain ingredients derived from GMO (genetically modified organisms) origin.



**VOERMOL**