

## PRODUCTION SUPPLEMENTS

### VOERMOL PRODUCTION LICK

Class: Protein, Mineral and Energy Supplement for Ruminants

**VOERMOL PRODUCTION LICK** is a complete, ready-mixed lick that provides protein, energy, minerals, vitamins and trace minerals for the optimum production of the grazing animal:

- contains grains and molasses as energy sources;
- helps to prevent excessive mass loss in cows after calving;
- helps to prevent pregnancy toxæmia in ewes during late gestation and can also be used for flush feeding of ewes;
- ensures weight gain of young growing animals on good quality veld;
- contains medicaments which promote energy usage from pastures.



#### COMPOSITION OF VOERMOL PRODUCTION LICK

REG NO V 10108 (Act 36/1947)

		(g/kg)			(mg/kg)
Crude Protein*	(min)	250	Manganese		100
Urea	(max)	55	Copper		40
Crude Fibre	(max)	80	Cobalt		0.8
Moisture	(max)	160	Iron		150
Calcium	(min/max)	10/12	Iodine		4
Phosphorus	(min)	8	Zinc		150
Potassium		18	Selenium		0.9
Magnesium	(min)	4	Vitamin A (IU/kg)		10 000
Sulphur	(min)	4	Energy estimate (MJ ME/kg)		8
*67.4 % derived from NPN			Contains medicaments		

Packaging – 50 kg

**See urea warning**

#### FEEDING INSTRUCTIONS

Use **Voermol Production Lick** as a protein and energy supplement to hay and/or veld.

#### RECOMMENDED INTAKE

(g/animal/day)

Cattle: 1 500 – 1 620\*

Sheep: 250 – 270\*

\*Maximum allowable intake of lick.

**NB. Contains medicaments and must therefore not be fed to horses.**



**VOERMOL**

## UREA WARNING

### UREA

The majority of licks contain urea that can be dangerous under certain conditions. During the formulation and manufacturing process of **Voermol** licks, great care is taken to minimise the risks of feeding urea. The possibilities of urea poisoning can be further reduced by taking note of the following:

- use licks as prescribed;
- limit lick intake to the prescribed maximum levels;
- feed the licks together with sufficient grazing and/or roughage;
- take special care during feed scarcity (drought conditions) to prevent high lick intake;
- prevent excessive lick intake in the initial stages (salt hunger) by allowing the animals access to **Voermol Rumevite 6P** for the first 14 days;
- do not let licks get wet.

**See urea warning on the back of the bag.**

### UREA WARNING

This farm feed contains NPN (Non-Protein Nitrogen) sources and must therefore be fed strictly according to instructions for use.

- Vinegar is an effective remedy against NPN poisoning. Mix with an equal amount of water. Dose half a bottle per calf/sheep or 2 – 4 bottles per head of cattle (1 bottle = 750 ml).
- Protect this farm feed against rain. NPN is soluble and animals drinking such a solution could be poisoned.
- Do not feed this farm feed indiscriminately with other NPN containing farm feeds. Consult an animal scientist.
- Mix concentrate thoroughly with the prescribed ingredients.
- This is a supplement and not a feed. Sufficient grazing and/or roughage must be available at all times.
- Keep lick troughs filled and prevent gluttonous eating by hungry animals. A constant daily intake can help prevent poisoning.
- Before feeding a NPN containing lick, feed an ordinary salt / phosphate lick for at least 7 days.

### GMO STATEMENT

These animal feeds may contain ingredients derived from GMO (genetically modified organisms) origin.



**VOERMOL**